A Clinical Prediction Model of Asymptomatic Left Ventricular Dysfunction in Survivors of Childhood Cancer in Ontario, Canada

Sickhids

POGG

PEDIATRIC ONCOLOGY GROUP OF ONTARIO

Roaa Shoukry ¹, Aasthaa Bansal ², Sameera Ahmed ³, Valerie Liu ³, Danielle Weidman ^{1, 3}, Paaladinesh Thavendiranathan ³, David Hodgson ^{3, 4}, Petros Pechlivanoglou ¹

1.The Hospital for Sick Children, Toronto, Canada; 2. University of Washington, USA; 3. University Health Network, Toronto, Canada; 4. Pediatric Oncology Group of Ontario, Toronto, Canada

QUHN

Background

- Survivors of childhood cancer (CCS) treated with anthracyclines or radiation have a higher risk of cardiomyopathy (CMP), like asymptomatic left ventricular dysfunction (ALVD) and heart failure (HF).
- Current guidelines recommend a risk-stratified schedule of screening. However, CCS
 may benefit from personalized prognostic screening for early intervention.
- A clinical prediction model which identifies CCS at high risk for ALVD can be used to determine different thresholds for continued screening. This may support precise decision-making to improve the health outcomes of CCS.

Objectives

- To develop a dynamic prediction model of ALVD risk using individual-level data in Ontario, Canada.
- To assess the impact of ejection fraction (EF) percentage on predicting ALVD risk

Methods

Data Sources: Retrospective chart data from the Princess Margaret Cancer Centre in Ontario

Eligibility Criteria: CCS treated with anthracyclines or radiation between 0-20 years who received at least two echocardiograms between 1980/01/01 to 2022/05/31

Outcome: ALVD, defined by ejection fraction (EF) < 51% (males); < 53% (females)

Time Horizon of Predictions: Two-year risk (prediction window) over 20 years (landmark time horizon)

Model: Landmark analysis with a multivariable Cox proportional hazards model **Assessments of Model Performance:** Calibration plots; time-varying AUC scores

Table 1. List of predictors for each prediction model

Model 1	Model 2 Age		
Age			
Age at diagnosis	Age at diagnosis		
Time between diagnosis and 1st screening	Time between diagnosis and 1st screening		
Time between screening and LM	Time between screening and LM		
No. of past screenings	No. of past screenings		
CMP risk high (vs not high)	CMP risk high (vs not high)		
EF percentage	Baseline EF		
Linear interactions with LM	Linear interactions with LM		
Quadratic interactions with LM	Quadratic interactions with LM		

Results


Table 2. Number of participants and events for each model

	Model 1: EF	Model 2: Baseline EF
N participants	691	691
N events	94	99
EF = ejection fraction		

Table 3. Sample characteristics

Mean age at diagnosis (Range)	7.44 (3 – 20)	CMP Risk (%)	
Mean attained age (Range)	31.11 (19 - 67)		
Sex: Female (%)	341 (49.4)	High	402 (58.2)
Treatment Exposure (%)		Moderate	189 (27.4)
Anthracycline	651 (94.2)	Moderate	100 (27.1)
Radiation	354 (51.2)	Low	100 (14.5)
		- CMP = cardiomyopathy	

Figure 1. (A) Calibration plots at times 0 to 10 yrs and (B) AUC scores for each model



Conclusion

This prediction model demonstrated high predictive performance of ALVD in CCS using standard clinical parameters. Both models perform similarly. The models may be used for precise medical decision-making in survivors of cancer. The time-varying hazard ratios generated will be used in the microsimulation-based cost-utility analysis for routine echocardiographic screening in survivors in Canada.