

The impact of clinical heterogeneity on conducting network meta-analyses for overall survival of previously untreated intermediate- or poor-risk advanced/metastatic renal cell carcinoma patients

Szabo N¹, Grevinga M¹, Malcolm B², Kurt M³, Ejzykowicz F³, Chun DS³, May JR², Kroep S¹

¹Pharmerit - an OPEN Health Company, Rotterdam, the Netherlands; ²Bristol Myers Squibb, Uxbridge, UK; ³Bristol Myers Squibb, Princeton, NJ, USA.

Background

- Kidney cancer, of which renal cell carcinoma (RCC) accounts for approximately 85%, is the 7th most common cancer worldwide in men, and the 10th most common cancer worldwide in women.¹
- Nivolumab (Opdivo®) is an immunoglobulin G4 human monoclonal antibody (IgG4 HuMAb) that binds to the programmed cell death-1 (PD-1) receptor, blocking the interaction of PD-1 with its ligands, PD-L1 and PD-L2.^{2,3}
- Knowledge concerning the comparability of clinical efficacy across interventions is essential beyond the available head-to-head comparisons, which would mostly include sunitinib as a comparator. A network meta-analysis (NMA) can be used synthesize evidence for differences in relative treatments; however, the validity of performing a NMA needs to be assessed by analyzing the networks of evidence and the heterogeneity across relevant trials.

Objective

- The current study investigates the feasibility of conducting a NMA for overall survival (OS) in the intermediate-/poor-risk population previously untreated aRCC patients.

Methods

- A systematic literature review (SLR) identified all published RCTs in previously untreated aRCC.⁴ Available evidence was synthesized by evaluating whether the pre-defined relevant interventions formed a network of evidence for OS outcomes in the intermediate-/poor-risk population.
- Clinical heterogeneity was assessed for each population, intervention, comparators, outcomes, and study design (PICOS):
 - Population:** age, sex, Eastern Co-operative Oncology Group performance status (ECOG-PS), Memorial Sloan Kettering Cancer Center score (MSKCC) and/or International Metastatic RCC Database Consortium score (IMDC), prior nephrectomy, prior use of radiation therapy, PD-L1 status, metastatic sites, race, region
 - Intervention:** treatment type, dose, and regimen
 - Outcomes:** definition of OS, stratified versus unstratified results
 - Study characteristics:** study phase, number of patients, study aim, study design (for example, cross-over design), follow-up duration
- Feasibility assessment was based on the framework by Cope et al. (2014).⁵
- The network of evidence was clustered based on eight relevant comparator treatment arms:
 - Avelumab + axitinib (AVE+AXI)
 - Cabozantinib (CAB)
 - Bevacizumab + interferon alfa (BEV+IFN)
 - Nivolumab + Ipilimumab (NIV+IPI)
 - Pazopanib (PAZ)
 - Pembrolizumab + axitinib (PEM+AXI)
 - Sunitinib (SUN)
 - Temsirolimus (TEM)

Results

Systematic Literature Review

- The SLR was performed and identified all available RCTs in previously untreated aRCC using MEDLINE, MEDLINE-IN-PROCESS, EMBASE, and the Cochrane library, the last update was on June 4th, 2020. A total of 14,027 records were identified, of which 121 satisfied the PICOS criteria. For the NMA, only RCTs were considered (N=57).⁴

Network Diagram

- For OS in the intermediate-/poor-risk population base case, the network included nine studies (see Table 1), which were relevant for forming a linked network. Table 2 reports the efficacy outcomes reported in the nine trials included. Finally, Figure 1 shows the network of evidence for OS in the intermediate/poor-risk population.

Table 1. Overview of the study characteristics and treatments of the trials included in the NMA				
Trial Name	Treatment	n	Study Phase	Study Design*
AVOREN ⁶	BEV+IFN	327	Phase 3	RCT
	IFN	322		
CABOSUN ^{7,8}	CAB	79	Phase 2	RCT
	SUN	78		
CALGB 90206 ^{9,10}	BEV+IFN	369	Phase 3	RCT
	IFN	363		
CheckMate-214 ¹¹	NIV+IPI	425	Phase 3	RCT
	SUN	422		
COMPARZ ¹²	PAZ	557	Phase 3	RCT
	SUN	553		
GLOBAL ARCC ¹³	TEM	209	Phase 3	RCT
	IFN	207		
JAVELIN Renal 101 ¹⁴	AVE+AXI	442	Phase 3	RCT
	SUN	444		
KEYNOTE-426 ¹⁵	PEM+AXI	432	Phase 3	RCT
	SUN	429		
NCT00083889 ¹⁶	SUN	375	Phase 3	RCT
	IFN	375		
TemPa ¹	TEM	35	Phase 2	RCT
	PAZ	34		

*Possible study design types: RCT, cross-over design and treatment sequencing

Figure 1. Network diagram for the intermediate-/poor-risk population




Table 2. Overview of the efficacy data of trials included in intermediate-/poor-risk population NMA			
Trial name	Treatment / Comparator	Risk population	OS HR (95% CI)
AVOREN ¹⁷	BEV+IFN / IFN	Intermediate	0.74 (0.53-1.02)
		Poor	0.87 (0.48-1.56)
CABOSUN ^{7,19}	CAB / SUN	Intermediate/poor	0.80 (0.53-1.21)
		Intermediate	0.87 (0.71-1.06)
CALGB 90206 ^{9,10}	BEV+IFN / IFN	Intermediate	0.75 (0.46-1.22)
		Intermediate/poor	0.66 (0.54-0.82)
CheckMate-214 ¹⁸	NIV+IPI / SUN	Intermediate	0.66 (0.50-0.87)
		Intermediate	0.66 (0.50-0.87)
		Poor	0.57 (0.39-0.82)
COMPARZ ¹²	PEM / SUN	Intermediate	0.90 (0.74-1.09)
		Intermediate	0.85 (0.56-1.28)
GLOBAL ARCC ¹³	TEM / IFN	Intermediate/poor	0.73 (0.58-0.92)
		Intermediate/poor	0.63 (0.50-0.71)
KEYNOTE-426 ^{15,19}	PEM+AXI / SUN	Intermediate	0.53 (0.35-0.82)
		Intermediate	0.53 (0.35-0.82)
		Poor	0.43 (0.23-0.81)
JAVELIN Renal 101 ¹⁴	AVE+AXI / SUN	Intermediate	0.86 (0.62-1.20)
		Poor	0.57 (0.36-0.90)
NCT00083889 ¹⁶	SUN / IFN	Intermediate	0.79 (0.62-1.00)
		Poor	0.66 (0.36-1.21)
TemPa ¹	TEM / PAZ	Intermediate/poor	1.16 (0.70-1.93)

Results (Continued)

Heterogeneity Assessment

- The heterogeneity assessment for the intermediate-/poor-risk population for OS included nine trials.
- Reporting was scarce for the baseline characteristics in the intermediate-/poor-risk population, and many of the trials could not be included due to a shortage of information.
- For the characteristics, such as age and sex, that were reported across a sufficient number of trials, populations appeared to be homogenous.


ECOG-PS

- ECOG-PS was only reported for two of the trials. Trials that reported on ECOG-PS scores have a large difference in distribution of scores between trials, in particular those with an ECOG-PS score of zero where the proportion ranged from 2.9% to 46% (Table 3).

Table 3. Distribution of ECOG-PS in studies included in the NMA					
Trial Name	Treatment	n	ECOG 0	ECOG 1	ECOG 2
CABOSUN ^{7,8}	CAB	79	46.0%	42.0%	13.0%
	SUN	78	46.0%	41.0%	13.0%
TemPa ¹	TEM	35	2.9%	40.0%	57.1%
	PAZ	34	2.9%	35.3%	61.8%


MSKCC/IMDC risk score

- Both the risk score definition as well as the distribution differed across trials. Figure 2 shows that even between studies that use the same definition, the risk score distribution varies (e.g. TemPa trial compared other studies).



Prior nephrectomy

- The percentage of patients who had a prior nephrectomy was similar (66% versus 80.2%). Only in the TemPa trial, the number of patients undergone a prior nephrectomy appears to be much lower in each treatment arm compared to the other trials (Figure 3).



Prior use of radiation therapy, PD-L1 expression, and metastatic sites

- Prior use of radiation therapy and PD-L1 expression was exclusively reported in the Checkmate 214 trial. Metastatic sites information, in particular bone metastasis, were reported across two studies only, namely Checkmate 214 and CABOSUN trials. However, the percentages of patients with bone metastasis differed between the two trials, ranging from 22% to 23% in the Checkmate 214 trial and 36% to 37% in the CABOSUN trial.

Race of patients

- There is homogeneity with regards to race. In both trials, the patients were predominantly white (>75%) (Table 4).

Table 4. Distribution of race of patients in studies included in the NMA						
Trial name	Treatment	n	White	Black	Asian	Other
CABOSUN ⁸	CAB	79	89.0%	4.0%	-	9.0%
	SUN	78	96.0%	3.0%	-	1.0%
TemPa ¹	TEM	35	77.1%	-	-	22.9%
	PAZ	34	76.5%	-	-	23.5%

Conclusions

- Due to the scarcity of reporting of relevant baseline characteristics for the intermediate-/poor-risk population, we can conclude that for several characteristics, such as prior use of radiation, PD-L1 expression, and metastatic sites, homogeneity across trials cannot be confirmed.
- For some of the characteristics that were reported across trials, such as age and sex, there seems to be homogeneity. However, differences were seen regarding the proportion of patients with higher ECOG-PS scores. Moreover, due to the unavailability of evidence we are unable to draw any conclusions for other characteristics, such as metastatic sites, PD-L1 expression, prior radiation, and race.
- In order to perform a NMA, we advise to limit heterogeneity in the evidence caused by prognostic risk score. We recommend excluding studies reporting outcomes only for the intermediate-risk population, and employing meta-regression to adjust for imbalances across all trials, where possible.

References

1. Tannir, N. M. et al. Temsirolimus versus Pazopanib (TemPa) in Patients with Advanced Clear-cell Renal Cell Carcinoma and Poor-risk Features: A Randomized Phase II Trial. *Eur Urol Oncol*, doi:10.1016/j.euo.2019.06.004 (2019).

2. George, S. et al. Safety and Efficacy of Nivolumab in Patients With Metastatic Renal Cell Carcinoma Treated Beyond Progression: A Subgroup Analysis of a Randomized Clinical Trial. *JAMA oncology* 2, 1179-1186, doi:10.1001/jamaoncol.2016.0775 (2016).

3. Escudier, B. et al. Treatment Beyond Progression in Patients with Advanced Renal Cell Carcinoma Treated with Nivolumab in CheckMate 025. *Eur Urol* 72, 368-376, doi:10.1016/j.eururo.2017.03.037 (2017).

4. Kram, C. W. et al. Efficacy and safety in previously untreated, advanced/metastatic renal cell carcinoma - A systematic literature review update. *Ispr Europ J Med* 2020 (2020).

5. Cope, S. et al. A process for assessing the feasibility of a network meta-analysis: a case study of everolimus in combination with hormonal therapy versus chemotherapy for advanced breast cancer. *BMC medicine* 12, 93 (2014).

6. Escudier, B. et al. Phase III trial of bevacizumab plus interferon alfa-2a in patients with metastatic renal cell carcinoma (AVOREN): final analysis of overall survival. *J Clin Oncol* 28, 2144-2150, doi:10.1200/JCO.2009.26.7849 (2010).

7. George, D. J. et al. Cabozantinib Versus Sunitinib for Untreated Patients with Advanced Renal Cell Carcinoma of Intermediate or Poor Risk: Subgroup Analysis of the Alliance A031203 CABOSUN Trial. *Oncologist* 24, 1497-1501, doi:10.1634/theoncologist.2019-0316 (2019).

8. Choueiri, T. K. et al. Cabozantinib Versus Sunitinib As Initial Targeted Therapy for Patients With Metastatic Renal Cell Carcinoma of Poor or Intermediate Risk: The Alliance A031203 CABOSUN Trial. *J Clin Oncol* 35, 591-597, doi:10.1200/JCO.2016.70.7398 (2017).

9. Rini, B. J. et al. Bevacizumab plus interferon alfa compared with interferon alfa monotherapy in patients with metastatic renal cell carcinoma: CALGB 90206. *J Clin Oncol* 26, 5422-5428, doi:10.1200/JCO.2008.16.9847 (2008).

10. Rini, B. J. et al. Phase III trial of bevacizumab plus interferon alfa versus interferon alfa monotherapy in patients with metastatic renal cell carcinoma: final results of CALGB 90206. *J Clin Oncol* 28, 2137-2143, doi:10.1200/JCO.2009.26.5561 (2010).

11. Motzer, Robert J., et al. "Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma." *New England Journal of Medicine* (2019).

12. Motzer, R. J., Hutson, T. E., McCarroll, L., Deen, K. & Choueiri, T. K. Overall survival in renal-cell carcinoma with pazopanib versus sunitinib. *N Engl J Med* 370, 1769-1770, doi:10.1056/NEJMc1400731 (2014).

13. Hudes, G. et al. Temsirolimus, interferon alfa, or both for advanced renal-cell carcinoma. *N Engl J Med* 356, 2271-2281, doi:10.1056/NEJMc1406838 (2007).

14. Motzer, R. J. et al. Avelumab plus axitinib versus sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med* 380, 1103-1115, doi:10.1056/NEJMc1816467 (2019).

15. Rini, B. J. et al. Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med* 380, 1116-1127, doi:10.1056/NEJMc1816714 (2019).

16. Motzer, R. J. et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol* 27, 3584-3590, doi:10.1200/JCO.2008.20.1929 (2009).

17. Escudier, B. et al. Bevacizumab plus interferon alfa-2a for treatment of metastatic renal cell carcinoma: a randomised, double-blind phase III trial. *The Lancet* 370, 2103-2111 (2007).

18. Tannir NM, et al. Thirty-month follow-up of the phase III CheckMate 214 trial of first-line nivolumab + ipilimumab (N+I) or sunitinib (S) in patients (pts) with advanced renal cell carcinoma (aRCC). *Journal of Clinical Oncology*. 2019;37(7 suppl):457-457. doi:10.1200/JCO.2019.37.7 suppl.454.

19. Plimack, E. R. et al. Pembrolizumab plus axitinib versus sunitinib as first-line therapy for advanced renal cell carcinoma (RCC): Updated analysis of KEYNOTE-426. *Journal of Clinical Oncology* 38, 5001-5001, doi:10.1200/JCO.2020.38.15, suppl. 5001 (2020).

Acknowledgments

- This study was supported by Bristol-Myers Squibb.