

BACKGROUND AND OBJECTIVES

Lung cancer represents the most frequent and lethal cancer worldwide [1]. Non-small cell lung cancer (NSCLC) is the predominant form of lung cancer [2], with about half of the cases diagnosed with Stage I-III disease. In spite of the advances in diagnostic procedures and the current therapeutic options, the management of NSCLC represents an open challenge for researchers and clinicians, above all for the high frequency of relapse which involve up to 60% of patients [3].

This real-world analysis aims to estimate the epidemiology of early-stage NSCLC (eNSCLC), patients' characteristics, and economic burden in a clinical practice setting in Italy.

REFERENCES


1. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, Forman D, Bray F. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. Eur J Cancer. 2013 Apr;49(6):1374-403. doi: 10.1016/j.ejca.2012.12.027.

2. Vansteenkiste J, Wauters E, Reymen B, Ackermann CJ, Peters S, De Ruyscher D. Current status of immune checkpoint inhibition in early-stage NSCLC. Ann Oncol. 2019 Aug 1;30(8):1244-1253. doi: 10.1093/annonc/mdz175.

3. Postmus PE. Changing health care costs for NSCLC, what does it mean? Lung Cancer. 2018 Mar;117:62-63. doi: 10.1016/j.lungcan.2017.11.006.

METHODOLOGY

An observational analysis was performed using administrative databases (ADB) linked to pathological anatomy-data (APD), covering around **2.5 mln health-assisted individuals**. From **2015-mid 2021** (study-period) eNSCLC patients (stage II-IIIa) were identified by the algorithm **Figure 1**.




INDEX-DATE: lung surgery-date (patients were characterized during 12-months before index-date and followed-up from index-date until study-period ending). Patients were stratified into those presenting **loco-regional or metastatic recurrence** during follow-up, and **annualized healthcare direct-costs** covered by the Italian National Health System (INHS) were estimated.

RESULTS


PREVALENCE AND INCIDENCE OF eNSCLC PATIENTS DURING 2019-2020 AND PROJECTION ON ITALIAN POPULATION

- On 2019-2020, the prevalence of eNSCLC was 104.3-117.1/million health-assisted subjects, and the annual incidence was 38.6-30.3/million (**Figure 2A**).
- Data projected to **Italian population** estimated 6,206 (2019) and 6,967(2020) prevalent and 2,297 (2019) and 1,803 (2020) incident cases (**Figure 2B**).



RECURRENCE AND ANNUALIZED HEALTHCARE DIRECT COSTS DURING FIRST 2 YEARS AFTER SURGERY AMONG ALIVE PATIENTS

- Overall, **458 eNSCLC patients were included**, aged 67.4±8.4 years (60.5% males).
- During 5-year follow-up period after surgery, **52.4%** of patients had a recurrence after a **median period of 20.5** (95%CI: 14.6-29.5) months (**not-shown**). At **24-months after surgery**, **45.6%** (N=99/217) **patients had recurrence** [6% (N=13/217) loco-regional and 39.6% (N=86/217) metastatic-recurrence] (**Figure 3**).
- Healthcare total direct-costs/patients averaged 11,714€ in no-recurrence and 21,577€ and 24,105€ in loco-regional and metastatic-recurrence patients, respectively (**Figure 3**).
- The cost related to drugs for lung cancer averaged **1,132€ in no-recurrence** and **4,307€ and 9,227€ in loco-regional and metastatic-recurrence patients**, respectively (**Figure 3**). Moreover, those for **visits/tests and lung cancer hospitalizations** averaged, respectively, 3,561€ and 5,260 € (in no-recurrence), 6,511€ and 7,041€ (in loco-regional recurrence), and 5,175€ and 6,240€ (in metastatic recurrence) per patients (**Figure 3**).



AUTHORS

VALENTINA PERRONE¹, ELISA GIACOMINI¹, DIEGO SANGIORGI¹, MARGHERITA ANDRETTA², FAUSTO BARTOLINI³, GIUSEPPE TAURINO⁴, MARCO BELFIORE⁵, EMILIA SICARI⁵, LUCA DEGLI ESPOSTI¹

1. CliCon S.r.l. Società Benefit, Health, Economics & Outcomes Research, 40121 Bologna, Italy;

2. UOC Assistenza Farmaceutica Territoriale, ULSS 8 BERICA, Vicenza, Italy;

3. Dipartimento Farmaceutico USL Umbria 2, Terni, Italy;

4. Dipartimento del farmaco Azienda USL Toscana Nord Ovest, Pisa, Italy;

5. Roche Spa, Monza, Italy

CONCLUSIONS

This **real-world analysis** provides insights into the epidemiology of eNSCLC, patients' characteristics, and economic burden, in Italy. About **45% of patients had a recurrence**, and in recurrence patients total direct costs covered by INHS **were almost 2-fold** those of no-recurrence patients. The results of the present analysis could suggest that for the management of eNSCLC an unmet clinical need remains, and therapeutic optimization at early stages is required to improve the disease's clinical and economic burden.