



Cell-based treatment increases Quality of Life and reduces amount of Knee Replacement Surgeries compared to current standard of care for knee osteoarthritis patients




Panaxea

Iris W.A. Boot¹, Georgina Shaw², Mary Murphy², Hubertus J.M. Vrijhoef¹

¹ Panaxea b.v., Den Bosch, Brabant, The Netherlands; ✉ iris.boot@panaxea.eu

² University of Galway, Galway, Ireland

ISPOR code **CO126**



Background

Osteoarthritis (OA) is a leading cause of disability worldwide, resulting in pain, structural changes in the bone and joint space, and limitation of motion [1]. Current treatments are designed to reduce pain and improve the mobility of joints instead of promoting the regeneration of damaged articular cartilage [2]. Cell and cell-based therapies are being developed for the prevention and treatment of OA [2]. This study was set up to assess the effectiveness of new cell-based therapies compared to current standard of care for patients with knee OA in the Netherlands.

Methods

A Markov model with 8 health states was developed to calculate the difference in health outcomes (i.e., quality adjusted life year (QALYs)), amount of total knee replacement (TKR) surgery and TKR revision (TKRR) surgery over 40 years. Knee OA health states were based on the Kellgren Lawrence (KL) scale. The eight health states are Healthy, KL I, KL II, KL III, KLIV, TKR, TKRR and Death. Input parameters were derived from literature [3, 5-10]. QALYs were discounted [4].




Figure 1: Model structure

Results

The average discounted QALYs per person over 40 years were 22.8 for patients receiving cell-based treatment, and 20.4 for patients receiving standard of care. The amount of TKR surgeries were 559 and 565, respectively, the amount of TKRR surgeries 21 and 28, respectively.

Conclusion

Based on the available evidence, the model shows that cell-based therapies could increase the QALYs of Dutch knee OA patients and could decrease the amount of TKR and TKRR surgeries. Effectiveness data of the cell-based treatment should be updated when more data becomes available. Future cost-effectiveness studies could be performed to also assess the cost-effectiveness.

AutoCRAT

Automated Cellular Robot-Assisted Technologies for translation of discovery-led research in Osteoarthritis (AutoCRAT) is a Horizon2020 project focused on delivering new cell and cell-based therapies for OA and joint repair using sustainable sources of cells and developing a closed, scalable and regulatory-compliant automated system to enable future production of therapeutic products.

References

- [1] Ferket, B.S., et al., *Impact of total knee replacement practice: cost effectiveness analysis of data from the Osteoarthritis Initiative*. *Bmj*, 2017. **356**: p. j1131.
- [2] Zhu, C., et al., *Mesenchymal stem cells in osteoarthritis therapy: a review*. *Am J Transl Res*, 2021. **13**(2): p. 448-461.
- [3] van der Woude, J.A., et al., *Knee Joint Distraction Compared to Total Knee Arthroplasty for Treatment of End Stage Osteoarthritis: Simulating Long-Term Outcomes and Cost-Effectiveness*. *PLoS One*, 2016. **11**(5): p. e0155524.
- [4] Garattini, L. and A. Padula, *Dutch guidelines for economic evaluation: 'from good to better' in theory but further away from pharmaceuticals in practice?* *J R Soc Med*, 2017. **110**(3): p. 98-103.
- [5] Wesseling, J., Boers, M., Viergever, M. A., Hilberdink, W. K., Lafeber, F. P., Dekker, J., & Bijlsma, J. W. (2016). Cohort Profile: Cohort Hip and Cohort Knee (CHECK) study. *International journal of epidemiology*, **45**(1), 36–44.
- [6] <https://www.lroi-report.nl/knee/survival-tka/short-and-long-term-revision/#by-procedure-year>
- [7] <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/37360ned/table?fromstatweb>
- [8] Bedene, A., et al., (2019). Opioid prescription patterns and risk factors associated with opioid use in the Netherlands. *JAMA network open*, **2**(8), e1910223-e1910223.
- [9] Spekenbrink-Spooren, A., et al., (2018). Higher mid-term revision rates of posterior stabilized compared with cruciate retaining total knee arthroplasties: 133,841 cemented arthroplasties for osteoarthritis in the Netherlands in 2007-2016. *Acta orthopaedica*, **89**(6), 640–645.
- [10] Rassir R, Siersevelt IN, van Steenberg LN, Nolte PA. Is obesity associated with short-term revision after total knee arthroplasty? An analysis of 121,819 primary procedures from the Dutch Arthroplasty Register. *Knee*. 2020 Dec;**27**(6):1899-1906. doi: 10.1016/j.knee.2020.09.020. Epub 2020 Nov 18. PMID: 33220579.

Research leading to these results has received funding from the European Union's Horizon 2020 research and innovation program (under grant agreement No 874671 and 667932). The material presented and views expressed here are the responsibility of the author(s) only. The EU Commission takes no responsibility for any use made of the information set out.

