ENHANCED HEALTHCARE THROUGH THE ADOPTION OF MOBILE HEALTH APPS

J. BRANTES FERREIRA, T. A. CALDEIRA

¹Pontifical Catholic University of Rio de Janeiro, Rio de Janeiro, Brazil



INTRODUCTION

Mobile Health (m-Health) technologies revolutionize healthcare by overcoming time and geographical barriers, crucially aiding developing countries. **Despite offering cost-effective, evidence-based solutions, adoption rates are low**, especially among the elderly, with many users discontinuing use quickly.


OBJECTIVES

Leveraging on models like TAM and UTAUT, propose a model to better understand and enhance m-Health adoption.

METHOD

- **Cross-sectional Survey** with 512 adult Brazilian respondents
- ** Structural Equations Modeling (SEM)

** Use of established scales to measure Behavioral Intention, Attitude, Perceived Ease of Use, Perceived Usefulness, Self-Efficacy, and Trust



RESULTS

**p<0.000

Sample Healthcare Provider distribution:

- 17.6% National Health System (SUS),
- 75% Private Health Plan
- 7.4% Direct Pay to Service Providers

* Significant effects for these relationships:

- patient Self-Efficacy perceptions on Perceived Ease of Use of wearable technology for health purposes (0.813),
- patient **Trust** in the mobile app on their Perceived **Usefulness** of the app (0.647),
- the impacts of Perceived **Usefulness** on patient Attitude

 toward using mHealth apps (0.868).

* The model could explain:

- 86.8% of the variance in patient Attitude,
- 62.2% of the variance in patient Behavioral Intention to adopt mHealth apps for consultations.

CONCLUSIONS

*p<0.05

***** Several issues influence the adoption of m-Health apps:

- the degree of mobile app diffusion among the population,
- the availability of healthcare options,
- the impact of patient selfconfidence,
- patient trust in the use and adoption of mobile technologies,
- patient perception of mobile app usefulness.

***Implications for management:**

- diffusion strategies should emphasize utilitarian attributes,
- developers should focus on usefulness,
- client security concerns should be minimized both regarding their health and their privacy.

REFERENCES

Chong, A. Y., Blut, M., & Zheng, S. (2022). Factors influencing the acceptance of healthcare information technologies: A meta-analysis. Information & Management, 59(3), 103604.

Huang, J.-C. (2010). Remote health monitoring adoption model based on artificial neural networks. Expert Systems With Applications, 37(1), 307–314.

Khalifa, M., & Shen, K. N. (2008). Explaining the adoption of transactional B2C mobile commerce. Journal of Enterprise Information Management, 21(2), 110–124.

Lee, T. (2005). The Impact of Perceptions of Interactivity on Consumer Trust and Transaction Intentions in Mobile Commerce. Journal of Electronic Commerce Research, 6(3), 165–180.

Sun, Y., Wang, N., Guo, X., & Peng, Z. (2013). Understanding the Acceptance of Mobile Health Services: A Comparison and Integration of Alternative Models. Journal of Electronic Commerce Research, 14(2), 183–199.

Van Velsen, L., Tabak, M., & Hermens, H. (2017). Measuring patient trust in telemedicine services: Development of a survey instrument and its validation for an anticoagulation web-service. International Journal of Medical Informatics, 97, 52–58.

World Health Organization - WHO. (2021). Mobile Health for Oral Health: An implementation guide. WHO.