

INTRODUCTION

Internationally, caplacizumab has been included as part of the initial treatment of acquired thrombotic thrombocytopenic purpura (aTTP) along with plasma exchange and immunosuppression¹⁻⁶. In Colombia, caplacizumab was approved in January 2022.

OBJECTIVE


To determine the therapeutic qualification of caplacizumab in conjunction with plasma exchange and immunosuppression, compared with the pre-caplacizumab standard regimen (plasma exchange and immunosuppression), using the methodology of the local HTA agency (Colombian Institute of Technological Evaluation in Health)^{7,8}.

METHODS

- Qualification was performed following the modified Delphi technique with a panel of experts composed of four haemato-oncologists, a pharmaceutical chemist, and one patient.
- For the qualification, the results of effectiveness and safety obtained through a systematic review of the literature (statistical significance), therapeutic threshold values (clinical significance), and degree of acceptability (willingness to use the technology) were considered.
- The threshold and acceptability were previously established in a deliberative process with the same panel of experts.

POSTER HIGHLIGHT:

This is the first Sanofi experience applying the new effectiveness and safety Colombian guideline for HTA. Treatment using caplacizumab together with plasma exchange and immunosuppression was considered superior to plasma exchange and immunosuppression for the treatment of patients with aTTP.



CONCLUSIONS

Treatment using caplacizumab together with plasma exchange and immunosuppression was considered superior for the standard treatment of aTTP, since it showed clinically significant benefits in critical outcomes and has a safety profile which is not different from its comparator.

RESULTS

- Fourteen effectiveness and safety outcomes were submitted for the qualification process (Table 1).
- Caplacizumab displayed clinical significance for some effectiveness outcomes, was not considered inferior in terms of safety, and showed acceptability for its use.
- By consensus, the panel determined that the technology of interest is superior to the pre-caplacizumab standard treatment, in terms of treatment response and composite outcome (aTTP-related death, aTTP recurrence, or at least one major thromboembolic event), and is not different for other effectiveness and safety outcomes.
- In overall terms, the panel established that using caplacizumab is superior to the pre-caplacizumab standard treatment.

REFERENCES

- Picod A et al. Should all patients with immune-mediated thrombotic thrombocytopenic purpura receive caplacizumab? Journal of Thrombosis and Haemostasis. 2021 Jan 10;19(1):58–67.
- Volker LA et al. Real-world data confirm the effectiveness of caplacizumab in acquired thrombotic thrombocytopenic purpura. Blood Adv. 2020 Jul 7;4(13):3085.
- Dutt T et al. Real-world experience with caplacizumab in the management of acute TTP. Blood. 2021 Apr 1;137(13):1731–40.
- Coppo P et al. A regimen with caplacizumab, immunosuppression, and plasma exchange prevents unfavorable outcomes in immune-mediated TTP. Blood. 2021 Feb 11;137(6):733–42.
- Zheng XL et al. ISTH guidelines for treatment of thrombotic thrombocytopenic purpura. International Society on Thrombosis and Haemostasis. 2020.
- Mingot Castellano ME, Pascual Izquierdo C, et al. Recomendaciones para el abordaje clínico de pacientes con púrpura trombocitopénica trombótica. Med Clin (Barc). 2022 Jun 24;158(12):630.e1-630.e14.
- Estrada-Orozco K et al. Manual metodológico para la elaboración de evaluaciones de efectividad clínica, seguridad y validez diagnóstica de tecnologías en salud. Instituto de Evaluación Tecnológica en Salud. 2022.
- IETS. Manual para la solicitud y emisión de conceptos sobre las evaluaciones de tecnologías en salud realizadas por terceros. Instituto de Evaluación Tecnológica en Salud. 2021