# Real-World Reduction of Oral Corticosteroid Use In Patients Treated With Dupilumab and With Moderate-To-Severe Asthma

Ajinkya Pawar<sup>1</sup>, Michael S. Blaiss<sup>2</sup>, Brian D. Modena<sup>3</sup>, Zhixiao Wang<sup>4</sup>, Asif H Khan<sup>5</sup>, Lucia De Prado Gómez<sup>6</sup>, Nami Pandit-Abid<sup>7</sup>, Amr Radwan<sup>4</sup>, Juby A Jacob-Nara<sup>7</sup>

Sanofi, Cambridge, MA, USA; Medical College of Georgia at Augusta University, Augusta, GA, USA; Modena Allergy & Asthma, La Jolla, CA, USA; Regeneron Pharmaceuticals, Inc., Tarrytown, NY, USA;

Sanofi, Chilly-Mazarin, France; Sanofi, Madrid, Spain; Sanofi Bridgewater, NJ, USA

# BACKGROUND

- Patients with severe asthma are reported to use systemic corticosteroids
   (SCS) more frequently and at high dosages; they are at increased risk
   of corticosteroid-related adverse events<sup>1</sup>
   Characteristic measured
- Dupilumab efficacy in oral corticosteroid (OCS) sparing was demonstrated in phase 3 VENTURE (NCT02528214), with a reduction of OCS dose by 70% for patients treated with dupilumab who were glucocorticoid dependent<sup>2</sup>

### OBJECTIVE

This real-world analysis investigated the reduction in OCS use in patients treated with dupilumab, both overall cohort and with OCS-dependent moderate-to-severe asthma

# METHODS

- The retrospective single-arm study analyzed 780 patients with moderateto-severe asthma treated with dupilumab
- 141 OCS-dependent patients were included (defined as average OCS daily dose ≥ 5 mg/day prednisone equivalent during at least 60 days before dupilumab initiation)

# CONCLUSION

Treatment with dupilumab reduced the number of days with OCS use, cumulative dose of OCS and the proportion of patients who used OCS ≥ 90 days, both in OCS-dependent and overall cohort population.

# RESULTS

Table 1. Dupilumab reduced OCS in patients with moderate-to-severe asthma after a 12-month follow-up

Characteristic measured	Baseline	Follow-up	Relative difference, %	P value				
Any OCS use, n (%)								
OCS-dependent population $(N = 141)$	141 (100)	112 (79.4)	-20.6	< 0.0001				
Overall Cohort (N = 780)	664 (85.1)	434 (55.6)	-34.7	< 0.0001				
Cumulative OCS dose, mg (prednisone-equivalent doses), mean (SD)								
OCS-dependent population $(N = 141)$	3,485.5 (3,306.0)	2,323.0 (2,695.1)	-33.4	0.0096				
Overall Cohort (N = 780)	1,349.7 (1987.9)	1,153.1 (1928.6)	-14.6	0.0009				
Number of days OCS used, mean (SD)								
OCS-dependent population $(N = 141)$	139.8 (83.4)	105.4 (98.8)	-24.6	0.32				
Overall Cohort (N = 780)	50.4 (69.2)	48.1 (74.5)	-4.54	0.17				
OCS use for ≥ 30 days, n (%)								
OCS-dependent population $(N = 141)$	141 (100)	91 (64.5)	-35.5	< 0.0001				
Overall Cohort (N = 780)	294 (37.7)	162 (20.8)	-44.8	< 0.0001				
OCS use for ≥ 60 days, n (%)								
OCS-dependent population $(N = 141)$	133 (94.3)	68 (48.2)	-48.9	< 0.0001				
Overall Cohort (N = 780)	144 (18.5)	101 (12.9)	-30.3	< 0.0001				
OCS use for ≥ 90 days, n (%)								
OCS-dependent population $(N = 141)$	93 (66.0)	54 (38.3)	-42.0	< 0.0001				
Overall Cohort (N = 780)	99 (12.7)	72 (9.2)	-27.6	0.0007				
SD, standard deviation								

Table 2. Dupilumab reduced severe asthma exacerbations<sup>a</sup> in patients treated with dupilumab after 12-month follow-up

	Characteristic measured	Baseline	Follow-up	Absolute difference	Relative difference, %	P value
Experienced any severe asthma exacerbations, n (%)						
	OCS-dependent population ( $N = 141$ )	118 (83.7)	71 (50.4)	-33.3	-39.8	< 0.0001
	Overall Cohort (N = 780)	551 (70.6)	285 (36.5)	-34.1	-48.3	< 0.0001
	Number of severe asthma exacerbation	ons experienc	ed, mean (SD	)		
	OCS-dependent population (N = 141)	3.5 (2.5)	2.4 (1.9)	-1.11	-31.7	0.015
	Overall Cohort (N = 780)	2.5 (1.8)	2.0 (1.7)	-0.51	-20.3	0.0006
<sup>a</sup> An inpatient visit with asthma as primary, admitting, or discharging diagnosis, an ER visit for asthma as primary, admitting, or discharging diagnosis, or an acute course of SCS (oral or injectable conducted as Acute courses of OCS are a prescription of OCS ≤ 30-day supply and with an asthma diagnosis within 7 days of the asthma-related medial visit. SCS prescribed within 7 days will be considered as						· ·

Table 3. ICS usage in patients treated with dupilumab at baseline vs 12-month follow-up

ER, emergency room; SCS, systemic corticosteroid

Characteristic measured	Baseline	Follow-up	Absolute difference	Relative difference, %	P value
Low dose of ICS used, n (%)					
OCS-dependent population (N = 141)	34 (24.1)	27 (19.1)	-5	-20.8	0.21
Overall Cohort (N = 780)	172 (22.1)	123 (15.8)	-6.3	-28.51	< 0.0001
Medium dose of ICS used, n (%)					
OCS-dependent population (N = 141)	40 (28.4)	28 (19.1)	-8.5	-29.9	0.0227
Overall Cohort (N = 780)	149 (19.1)	106 (13.6)	-5.5	-28.8	< 0.0001
High dose of ICS used, n (%)					
OCS-dependent population (N = 141)	63 (44.7)	47 (33.3)	-11.4	-25.5	0.009
Overall Cohort (N = 780)	363 (46.5)	230 (29.5)	<b>–17</b>	-36.6	< 0.0001
ICS, inhaled corticosteroids					

References: 1. O'Byrne PM, et al. Am J Respir Crit Care Med. 2009;179:19-24. 2. Rabe KF, et al. N Engl J Med. 2018;378:2475-85.

Acknowledgments and funding sources: Research sponsored by Sanofi and Regeneron Pharmaceuticals Inc. ClinicalTrials.gov Identifier: NCT02528214. Medical writing/editorial assistance was provided by Bruno Manso, PhD, of Excerpta Medica, and was funded by Sanofi and Regeneron Pharmaceuticals Inc. according to the Good Publication Practice guideline.

Disclosures: Pawar A, Khan AH, de Prado Gómez L, Pandit-Abid N, Jacob-Nara JA: Sanofi – employees, may hold stock and/or stock options in the company. Blaiss MS: Amgen, AstraZeneca, Covis, Regeneron Pharmaceuticals Inc., Sanofi, Teva – honoraria, consulting fees. Radwan A, Wang Z: Regeneron Pharmaceuticals Inc. – employee and shareholder.

